



NEW ZEALAND BIOSECURE

Entomology Laboratory



## *Culex (Culex) annulirostris* (Skuse)

common banded mosquito

**NZ Status: Not present – Unwanted Organism**



### Vector and Pest Status

*Culex annulirostris* is a vector of Murray Valley encephalitis (Kay *et al.*, 1984; Lee *et al.*, 1989) Japanese encephalitis (van den Hurk *et al.*, 2003), Kunjin (Kay *et al.*, 1984), Ross River virus (Harley *et al.*, 2000) and Barmah Forest virus (Boyd and Kay, 2000). In most localities where Bancroftian filariasis (*Wuchereria bancrofti*) is endemic, *Cx. annulirostris* has little or no significance as a vector (Lee *et al.*, 1989). However, in Irian Jaya where nocturnal periodic filariasis is prevalent, it is a major vector (Lee *et al.*, 1989).

This species is also able to carry dog heartworm (Russell and Geary, 1992; 1996) and is probably a major vector of myxomatosis (Russell, 1993). Other viruses that have been associated with *Cx. annulirostris* include Alfuy, Edge Hill, Eubenangee, Gan Gan, Kokobera, Kowanyama, Rift Valley fever, Trubanaman, and Sindbis (Lee *et al.*, 1989).


### Geographic Distribution

This mosquito is found in Australia, parts of Southeast Asia and the Pacific, to the east through the island chains as far as French Polynesia, and north through Papua New Guinea to the Mañana Islands. It has been recorded from the Austral Islands,

Bismarck Archipelago, Cook Islands, Fiji, Indonesia, Kiribati, Moluccas, Nauru, New Caledonia, Palau, Philippines, Samoa, Society Islands, Rotuma Island, Solomon Islands, Tonga, Torres Strait Islands, Tuamotu Archipelago, Tuvalu, Vanuatu and Wallis and Futuna Islands (Lee *et al.*, 1989).

In Australia, it is found in New South Wales (widespread coastal and inland), Victoria (widespread but not common south of the Central Highlands), South Australia (widespread, particularly Murray Valley), Tasmania (east coast but only one record) as well as Queensland, the Northern Territory and Western Australia (Russell, 1993).

It is believed that *Cx. annulirostris* would be unable to establish in cooler parts of New Zealand.



This map denotes only the country or general areas where this species has been recorded, not actual distribution.

### Incursions and Interceptions

This species has been intercepted in New Zealand 11 times. The last interception in 2016 was in Auckland International Airport, a specimen was found alive together with two *Aedes vexans* and seven *Aedes vigilax* in relation to a tent (camping equipment) that was used in New Caledonia (NZBEL Interception records, 2023)

### Taxonomy

*Culex annulirostris* belongs in Group A, the *sitiens* subgroup of the subgenus *Culex* (Lee *et al.*, 1989). At present, this species is thought to be made up of five geographically restricted, divergent lineages, two of which are localised in Australia, two in Papua New Guinea, the Torres Strait and the top of Cape York, and the fifth identified from the Solomon Islands (Hemmenter *et al.*, 2006).

### Habits and Habitat

*Culex annulirostris* breeds in a variety of habitats. It is typically found in temporary freshwater pools on grassland as well as freshwater ponds, wetlands, swamps, lakes, dams, irrigated areas, drainage ditches, flooded rice fields and hoof prints (Lee *et al.*, 1989). It has also been collected in some artificial containers such as a 44 gallon drum, tyres, a stock drinking trough, canoes, a swimming pool and a bath (Lee *et al.*, 1989).

Kay *et al.* (1981) characterised typical larval habitats as fresh water swamps, lagoons and transient grassy pools, but also a variety of other sunlit or shaded habitats, usually with aquatic or emergent vegetation (Lee *et al.*, 1989). *Culex annulirostris* has also been found in waters of salinity up to 16.3 gm total salts/litre (Woodhill, 1936 in Lee *et al.*, 1989).

Eggs are laid in a boat shaped egg raft on the water surface, which usually consists of 50-200 eggs (Kay *et al.*, 1981). The eggs are not desiccation resistant and will die if they become dry (Kay *et al.*, 1981). In the laboratory, development from egg to adult ranged from 8.57 days at 35°C to 37 days at 15°C (McDonald *et al.*, 1980 in Lee *et al.*, 1989). Larval survival of this species under laboratory conditions was best at 25°C (McDonald *et al.*, 1980 in Lee *et al.*, 1989). In the field (in Australia), egg hatch to adult emergence took 21-32 days at water temperatures of 13-18°C (Kay *et al.*, 1981).

*Culex annulirostris* is a crepuscular species (Williams and Kokkinn, 2005) with adult females host feeding primarily at dusk and dawn, however this species will also feed during the day and at night, attacking both indoors and outdoors (Lee *et al.*, 1989). This species has been recorded feeding on humans, rabbits, dogs, fowl, cats, pigs, sheep, goats, horses, kangaroos, ox, buffalo cattle, chickens, guinea pigs, mice, brush-tailed possums, bats reptiles and amphibians (Lee *et al.*, 1989).

In a mark-release-recapture study in New South Wales, the maximum flight distance observed was 12km (the limit of the trapping network). The rate of dispersal of the population was estimated at 2.2km/day, with a mean distance travelled of 4.4km, however some of the population dispersed more than 5km (Bryan *et al.*, 1992).

## References

- Boyd, A.M. and Kay, B.H. 2000. Vector competence of *Aedes aegypti*, *Culex sitiens*, *Culex annulirostris* and *Culex quinquefasciatus* (Diptera: Culicidae) for Barmah Forest virus. *Journal of Medical Entomology* 37(5): 660-663.
- Bryan, J.H., O'Donnell, M.S., Berry, G. and Carvan, T. 1992. Dispersal of adult female *Culex annulirostris* in Griffith, New South Wales, Australia: a further study. *Journal of the American Mosquito Control Association* 8(4): 398-403.
- Derraik, J.G.B. 2004. Exotic mosquitoes in New Zealand: a review of species intercepted, their pathways and ports of entry. *Australian and New Zealand Journal of Public Health* 28(5): 433-444.
- Harley, D., S. Ritchie, D. Phillips, and A. van den Hurk. 2000. Mosquito isolates of Ross River virus from Cairns, Queensland, Australia. *American Journal of Tropical Medicine and Hygiene* 62:561-565.
- Hemmenter, S., Slapeta, J., van den Hurk, A.F., Cooper, R.D., Whelan, P., Russell, R., Johansen, C.A., and Beebe, N.W. 2006. Phylogeography of *Culex annulirostris* and *Culex palpalis*: arbovirus vector of Australia and Papua New Guinea. *7<sup>th</sup> Conference of the Mosquito Control Association of Australia*. 15-18<sup>th</sup> August, Novotel, Palm Cove, Queensland, Australia.
- Kay, B.H., Fanning, I.D. and J.G. Carley. 1984. The vector competence of Australian *Culex annulirostris* with Murray Valley Encephalitis and Kunjin viruses. *Australian Journal of Experimental Biology and Medical Science* 62(5): 641-650.
- Kay, B.H., Sinclair, P. and Marks, E.N. 1981. Mosquitoes: Their interrelationship with man. Pp. 157-174. In: *The ecology of pests – some Australian case histories*. (Ed.) Kitching, R.L. and Jones, R.E. CSIRO, Melbourne.

- Lee, D.J., Hicks, M.M., Debenham, M.L., Griffiths, M., Marks, E.N., Bryan, J.H. and Russell, R.C. 1989. *The Culicidae of the Australasian region*. Volume 7. Canberra, Australian Government Publishing Service.
- McDonald, G., McLaren, I.W., Shelden, G.P. and Smith, I.R. 1980. The effect of temperature on the population growth potential of *Culex annulirostris* Skuse (Diptera: Culicidae). *Australian Journal of Ecology* 5: 379-384.
- Russell, R.C. 1993. *Mosquitoes and mosquito-borne disease in southeastern Australia: a guide to the biology, relation to disease, surveillance, control and the identification of mosquitoes in southeastern Australia*. 2<sup>nd</sup> Edn. Sydney, University of Sydney pp. 318.
- Russell, R.C. and Geary, M.J. 1992. The susceptibility of the mosquitoes *Aedes notoscriptus* and *Culex annulirostris* to infection with dog heartworm *Dirofilaria immitis* and their vector efficiency. *Medical and Veterinary Entomology* 6: 154-158.
- Russell, R.C. and Geary, M.J. 1996. The influence of microfilarial density of dog heartworm *Dirofilaria immitis* on infection rate and survival of *Aedes notoscriptus* and *Culex annulirostris* from Australia. *Medical and Veterinary Entomology* 10(1): 29-34.
- Van den Hurk, A.F., Nisbet, D.J., Hall, R.A., Kay, B.H., MacKenzie, J.S. and Ritchie, S.A. 2003. Vector competence of Australian mosquitoes (Diptera: Culicidae) for Japanese encephalitis virus. *Journal of Medical Entomology* 40(1): 82-90.
- Williams, C.R. and Kokkinn, M.J. 2005. Daily patterns of locomotor and sugar-feeding activity of the mosquito *Culex annulirostris* from geographically isolated populations. *Physiological Entomology* 30: 309-316.
- Woodhill, A.R. 1936. Observations and experiments on *Aedes concolor*, Taylor (Diptera: Culicidae). *Bulletin of Entomological Research* 27: 633-648.