



NEW ZEALAND BioSECURE

Entomology Laboratory



## *Anopheles (Cellia) farauti* s.s. Laveran

**NZ Status: Not Present – NSP Watchlist**




### Vector and Pest Status

*Anopheles farauti* s.s. is a vector of malaria, and is highly susceptible to *Plasmodium vivax* and *P. falciparum* (Lee *et al.*, 1988). It is also a vector of *Bancroftian filariasis* (*Wuchereria bancrofti*) (Lee *et al.*, 1988).

### Geographic Distribution

The *Anopheles punctulatus* group (including *An. farauti*) is present in Australia, Indonesia, Solomon Islands, Papua New Guinea, Solomon Islands, Bismarck Archipelago and the islands of Vanuatu (Lee *et al.*, 1988). *Anopheles farauti* s.s. is the most widely

distributed species of the group and is found throughout northern Australia east to the islands of Vanuatu (Beebe *et al.*, 2000b).



This map denotes only the country or general areas where this species has been recorded, not actual distribution.

### Interceptions and Incursions

This species has not been intercepted in New Zealand.

### Taxonomy

*Anopheles farauti* is part of the *An. punctulatus* group, which are important vectors of malaria in the southwest Pacific. Presently the group is considered to contain about 12 cryptic species with overlapping morphology (Beebe *et al.*, 2000b), including

- *An. farauti* s.s. Laveran (formerly *An. farauti* No. 1) (Schmidt *et al.*, 2001)
- *An. hinesorum* Schmidt sp.n. (formerly *An. farauti* No. 2) (Schmidt *et al.*, 2001)
- *An. torresiensis* Schmidt sp.n. (formerly *An. farauti* No. 3) (Schmidt *et al.*, 2001)
- *An. farauti* No. 4-6,
- *An. irenicus* Schmidt sp.n. (formerly *An. farauti* No. 7) (Schmidt *et al.*, 2003)
- *An. punctulatus* Donitz
- *An. sp. near punctulatus*
- *An. koliensis* Owen
- *An. clowi* Rozeboom and Knight (Cooper *et al.*, 2000)

Some members of this group cannot be reliably identified by morphological markers, making field studies on their biology and behaviour difficult, only recently has work to determine their distributions been initiated (Beebe *et al.*, 2000b).

### Habits and Habitats

*Anopheles farauti* s.s. is a coastal species which has larvae that are tolerant of saline conditions (Sweeney, 1987), although they will also breed in fresh water (Beebe *et al.*, 2000b). Breeding sites include the margins of creeks and rivers entering the sea, pools and swamps formed behind beaches and coastal sand dunes (Beebe *et al.*, 2000b), semi-permanent ground waters with emergent vegetation such as swamps, lagoons and ponds, pig wallows, garden pools. Containers such as tins, coconut shells drums and canoes are an exception rather than the rule (Lee *et al.*, 1988).

This species is capable of crossing large expanses of water as indicated by its presence in the Solomon and Vanuatu islands (Beebe *et al.*, 2000b). Perry (1945) observed that females did not mate within two days of emergence.

In a biting study on the Solomon Islands investigating three members of the complex, *An. farauti* s.s. was the only species collected on human bait, with a much higher biting rate early in the evening between 6 30pm-8pm than later between 9pm-12am (Beebe et al., 2000a). Other hosts include dog, pig, horse, cat, cattle, goat, ruminant, possum, kangaroo. Adult females predominantly bite at night, both indoors and outdoors. Variations in peak biting activity have also been recorded (Lee *et al.*, 1988). Studies on dispersal have shown that females could fly 1 mile [1.6km] from the nearest larval habitat (F.H.S. Roberts & O'Sullivan, 1948) while the distance flown by marked females was generally less than 50m (Charlwood, Dagoro & Paru, 1985).

## References

- Beebe, N.W. and Saul, A. 1995. Discrimination of all members of the *Anopheles punctulatus* complex by polymerase chain-restriction fragment length polymorphism analysis. *American Journal of Tropical Medicine and Hygiene* 53(5) 478-481.
- Beebe, N.W., Bakote'e, B., Ellis, J.T. and Cooper, R.D. 2000a. Differential ecology of *Anopheles punctulatus* and three members of the *Anopheles farauti* complex of mosquitoes on Guadalcanal, Solomon Islands, identified by PCR-RFLP analysis. *Medical and Veterinary Entomology* 14(3) 308-312.
- Beebe, N.W., Cooper, R.D., Foley, D.H. and Ellis, J.T. 2000b. Populations of the south-west Pacific malaria vector *Anopheles farauti* s.s. revealed by ribosomal DNA transcribed spacer polymorphisms. *Heredity* 84 244-253.
- Charlwood, J.D., Dagoro, H. and Paru, R. 1985. Blood-feeding and resting behavior in the *Anopheles punctulatus* Donitz Complex (Diptera: Culicidae) from coastal Papua New Guinea. *Bulletin on Entomological Research* 75 463-475.
- Cooper, R.D., Waterson, D.G.E., Bangs, M.J. and Beebe, N.W. 2000. Rediscovery of *Anopheles (Celia) clowi* (Diptera: Culicidae), a rarely recorded member of the *Anopheles punctulatus* group. *Journal of Medical Entomology* 37(6) 840-845.
- Lee, D.J., Hicks, M.M., Griffiths, M., Debenham, M.L., Bryan, J.H., Russell, R.C., Geary, M. and Marks, E.N. 1987. The Culicidae of the Australian Region. Volume 5. *Entomology Monograph No. 2*. Canberra; Australian Government Publishing Services pp 315.
- Perry, W.J. 1945. Notes on the biology of the Malaria vector in the New Hebrides – Solomon Islands. *Malaria and Epidemic Disease Control*, pp. 16.
- Roberts, F. H. S. and O'Sullivan, P.J. 1948. Studies on the behaviour of adult Australasian Anophelines. *Bulletin of Entomological Research* 39, 159-178.
- Service, M.W. 2000. *Medical Entomology for Students*. 2<sup>nd</sup> edn. University Press, Cambridge, pp. 283.
- Schmidt, E.R., Foley, D.H., Hartel, G.F., Williams, G.M. and Bryan, J.H. 2001b. Descriptions of the *Anopheles (Celia) farauti* complex of sibling species (Diptera Culicidae) in Australia. *Bulletin of Entomological Research* 91(5) 389-411.
- Schmidt, E.R., Foley, D.H., Bugoro, H. and Bryan, J.H. 2003. A morphological study of the *Anopheles punctulatus* group (Diptera Culicidae) in the Solomon Islands, with a description of *Anopheles (Celia) irenicus* Schmidt, sp.n. *Bulletin of Entomological Research* 93(6) 515-526.
- Sweeney, A.W. 1987. Larval salinity tolerances of the sibling species of *Anopheles farauti*. *Journal of the American Mosquito Control Association* 3 589-592.